

**2003  
WHITLOCKS BAY SPAWNING STATION  
ANNUAL REPORT**

**Robert P. Hanten, Reservoir Biologist  
Pierre, South Dakota**

**South Dakota  
Department of  
Game, Fish and Parks  
Wildlife Division  
Joe Foss Building  
Pierre, South Dakota 57501-3182**

**Annual Report  
No. 04-01**

## **PREFACE**

This report discusses the 2003 season at Whitlocks Bay Spawning Station and previous data showing trends during the twenty years of station operation. The author would like to acknowledge all those who assisted with spawning operations, data collection, and maintenance of the facility. The author would also like to thank Sylvester Schied, Butch Schunot, Kyle Potter and Aaron Leingang for assisting with the coded-wire-tag (CWT) reading and station startup and shutdown.

### **Summary Table**

Summary of Spawn-Taking Effort Whitlocks Bay Spawning Station, 2003					
Species	Number returned	Females spawned	Eggs taken	Total cost of spawn	Cost Per 1000 eggs
Chinook salmon	459	284*	1,168,978	\$20,720	\$17.73

Rainbow trout were not included in this table due to the station not being operated during the spring.

\*Does not include green, spent, or bad females spawned

## Table of Contents

	Page
Preface and Summary Table.....	i
Table of Contents.....	ii
List of Tables.....	ii
List of Figures.....	iv
List of Appendices.....	v
Introduction.....	1
Rainbow Trout.....	4
Stocking.....	4
Station Returns and Egg Collection.....	4
Age and Growth.....	4
Chinook Salmon.....	5
Stocking.....	5
Station Returns and Egg Collection.....	5
Age and Growth.....	5
Contribution of North Dakota Salmon.....	6
Tri-State Chinook Salmon Egg Take and Inter-State Assistance.....	7
Visitation and Tours.....	10
Literature Cited.....	11

## **List of Tables**

## **Page**

1.	Costs of taking rainbow trout eggs at Whitlocks Bay Spawning Station, 2003.....	4
2.	Number of female Chinook salmon spawned and eggs collected, Whitlocks Bay Spawning Station, 2003.....	5
3.	Age composition, length and weight of 204 coded-wire tagged Chinook salmon collected at Whitlocks Bay Spawning Station, fall 2003.....	6
4.	Costs of taking Chinook salmon eggs at Whitlocks Bay Spawning Station, 2003.....	9

## List of Figures

	Page
1. Location of Whitlocks Bay Spawning Station.....	2
2. General layout of Whitlocks Bay Spawning Station.....	3
3. Chinook salmon returns to Whitlocks Bay Spawning Station compared to station intake water temperatures, Lake Oahe, SD, fall 2003.....	8

## List of Appendices

	Page
A. Rainbow trout stocked in Lake Oahe, 1996 - 2003.....	12
B. Numbers of coldwater species stocked in Lake Oahe, 1983 - 2003.....	13
C. Pounds and kilograms of coldwater species stocked in Lake Oahe, 1982 - 2003.....	14
D. Summary of rainbow trout return and egg collection at Whitlocks Bay Spawning Station, 1984 - 2003.....	15
E. Summary of salmon return and egg collection at Whitlocks Bay Spawning Station, 1984 - 2003.....	16
F. Chinook salmon stocked by number, size and mark in Lake Oahe, 1998 - 2003.....	17
G. South Dakota Chinook salmon coded-wire-tag data 1989 -2003.....	18
H. North Dakota Chinook salmon coded-wire-tag data 1993 -2003.....	19
I. South Dakota rainbow trout coded-wire tag data 1991 -2003.....	20
J. Number of coded-wire-tagged rainbow trout stocked in Lake Oahe, 1987 - 2003. Number stocked is corrected for tag retention.....	21
K. Number of coded-wire-tagged Chinook salmon stocked in Lake Oahe, 1987 - 2003. Number stocked is corrected for tag retention.....	22
L. Percent recovery of coded-wire tags returned by anglers and recovered at Whitlocks Bay Spawning Station in Lake Oahe 1988-2003.....	24
M. Salmon and trout cost breakdown explanation for each category in spring and fall budgets.....	25
N. Calculation of estimated contribution of North Dakota Chinook salmon to the spawn of Lake Oahe, South Dakota 2003.....	26
O. Summary of South Dakota, North Dakota and Montana Chinook salmon total egg take, distribution, or inter-state assistance.....	27
P. Length at selected number per pound values for rainbow trout and Chinook salmon.....	28
Q. Chinook salmon carcass contract results for 2003.....	29

## **Introduction**

In the early 1970's, the South Dakota Department of Game, Fish and Parks attempted to develop a salmonid sport fishery in Lake Oahe that would utilize the lake's coldwater habitat and diversify the existing fishery. Introductions of kokanee salmon, Bonneville cisco, opossum shrimp, lake herring and lake whitefish were made in hopes of establishing a forage base for a large predator species. These stockings, except for lake trout, were generally unsuccessful. Lake trout were selected because of the possibility that sufficient natural reproduction might occur to maintain a fishery without the need for annual maintenance stocking. However, minimal success was achieved with this species and no natural reproduction of this species was documented. In 1971, 7,500 adult Lake Superior rainbow smelt were stocked into Lake Sakakawea North Dakota and by 1976 smelt had established a self-sustaining population in downstream Lake Oahe. Chinook salmon had also reached Lake Oahe, as early as 1979, from Lake Sakakawea. As a result of the success of Lake Sakakawea rainbow smelt and Chinook salmon introductions in North Dakota, the South Dakota Department of Game, Fish and Parks implemented its own Chinook salmon program in 1982.

Chinook salmon from the Little Manistee Hatchery in Michigan and a hatchery in Wisconsin had been stocked as smolts into Lake Sakakawea, North Dakota in 1978. The origination of these Great Lakes Chinook were from two ocean-run strains from the state of Washington. The first was a Tule' strain Chinook which migrated up the Columbia River and spawned in the feeder streams of the Cascade Mountain Range at Spring Creek Hatchery. The second was a Puget Sound strain from Washington's Green River Hatchery. After three years of successful planting of smolts from these West Coast eggs, Michigan became self-sufficient for Chinook salmon production. By 1979 Chinook salmon stocked in Lake Sakakawea had reached Lake Oahe and by 1981 they were abundant enough that South Dakota Department of Game, Fish and Parks personnel collected 100,000 eggs from 54 female Chinook salmon at Whitlocks Bay, Lake Oahe, SD, October 19-22, 1981. Cleghorn Springs Hatchery incubated the eggs and produced 31,280 smolts which were stocked in Whitlocks Bay, April 1982. An additional 260,870 smolts produced from Lake Michigan eggs were also stocked in Lake Oahe. South Dakota has been relatively self-sufficient for salmon eggs since 1984.

The purpose of Whitlocks Bay Spawning Station (WBS) is to collect spawning adults of Chinook salmon and rainbow trout so that annual egg-production needs could be met, and a sport fishery in Lake Oahe could be maintained. Artificial propagation of Chinook salmon and rainbow trout is necessary because suitable spawning habitat is not available in Lake Oahe's warm and turbid incoming tributaries. The station has also been used to collect brown trout eggs; however, brown trout have not been stocked in Lake Oahe since 1990. Brown trout have not returned to the station since 1994.

Construction of Whitlocks Bay Spawning Station began in 1982 and the station was fully operational in April, 1984. The Whitlocks Bay Spawning Station consists of a fish ladder (artificial stream), four concrete holding ponds (45' X 8' X 4'), a crowding raceway, a 28' X 48' spawning building, and a water supply system (Figure 2). Two submersible pumps, capable of delivering a total of 2,600 gallons of water per minute (depending on lake elevation), are mounted on a skid that is pushed into the reservoir before the pumping season and removed when not in use. Water is pumped from the bay into the station where it gravity flows through raceways and down the fish ladder into the reservoir. The station is located 18 miles west of Gettysburg on Lake Oahe at Whitlocks Bay (Figure 1).

The station is operated from approximately October 7<sup>th</sup> through October 23<sup>rd</sup>. Personnel needed at the facility include a manager, a conservation technician and two seasonal employees. Considerable time is spent explaining department fisheries programs and providing guided tours to the public. In the fall, four to six additional people are needed during days of salmon egg collection.

This report discusses the 2003 season at Whitlocks Bay Spawning Station and previous data showing trends during the nineteen years of station operation. For a summary of previous operations, refer to annual reports for the years 1984-2002.




Figure 1. Location of Whitlocks Bay Spawning Station.






Figure 2. General layout of Whitlocks Bay Spawning Station

## **Rainbow Trout**

### **Stocking**

No rainbow trout were stocked in Lake Oahe in 2003 (Appendix B).

### **Station Returns and Egg Collection**

Whitlocks Bay Spawning Station was not operated this spring due to the overabundance of predators and extreme low abundance of prey fish. This current scenario calls for no stocking of rainbow trout. Past stocking records of rainbow trout for Lake Oahe are presented in Appendices A, B and C.

A summary of past rainbow trout returns and egg collection efforts can be found in Appendix D. No rainbow trout eggs were collected in 2003 therefore no cost was associated with categories listed in Table 1. An explanation of how the cost breakdown was completed for each category can be found in Appendix G.

**Table 1.** Costs of taking rainbow trout eggs at Whitlocks Bay Spawning Station, 2003.

---

Manpower/per diem/travel	\$0
Utilities (electricity and phone)	\$0
Misc.-building and grounds maintenance, hardware supplies etc.	<u>\$0</u>
<b>Total Costs</b>	\$0
Number of eggs taken	0
Cost/1,000 eggs collected	\$0

---

### **Age and Growth**

No rainbow trout coded wire tags were collected in 2003. Further breakdown of CWT returns, numbers of fish tagged, and tag retention rates can be found in Appendix I.

## **Chinook Salmon**

### **Stocking**

Seventeen thousand two hundred-fifty nine juvenile Chinook salmon were stocked in Lake Oahe in 2003. For a more detailed summary of current and past Chinook salmon stocking, refer to Appendix B, C, F and G.

### **Station Returns and Egg Collection**

On October 3, 2003 the two submersible pumps were turned on, producing a flow of approximately 2,600 gallons of water per minute down the ladder. Due to extremely low lake elevation (1578.8 [ftmsl.](#) by October 21<sup>st</sup>) a total of 34 metal culvert extensions were added on the end of the concrete ladder. Intake water temperatures decreased from 59° to 56° F during the pumping period. Fifteen Chinook salmon ascended the fish ladder the first night. Water temperature was 59° F on October 4<sup>th</sup>. The first salmon usually comes up the ladder the last week of September but due to the late start of the station the first salmon came up during the first week of October. Salmon returns produced bimodal peaks on October 9-11 and October 16-18 (Figure 3). Water temperature during this time remained fairly constant 57-59° F but occurred at warmer temperatures compared to the average water temperatures from 1998 thru 2002 of 53-55° F. Four hundred and fifty nine Chinook salmon ascended the ladder during the fall of 2003. Three salmon egg-take operations yielded approximately 1,168,978 eggs (Table 2). Two-hundred and eighty-four females returned via the fish ladder compared to 339, and 96 in 2002 and 2001 respectively. For a summary of past Chinook salmon returns, egg collection, and percent egg eye-up for WBS spawning operations, refer to Appendix E.

**Table 2.** Number of female Chinook salmon spawned and eggs collected, Whitlocks Bay Spawning Station, 2003.

<b>Date</b>	<b>Spawned</b>	<b>Green</b>	<b>Spent</b>	<b>Bad</b>	<b>Eggs collected</b>
10/8	69	3	0	3	299,500
10/14	110	0	1	13	442,575
10/21	105	0	0	11	426,903
<b>Total</b>	<b>284</b>	<b>3</b>	<b>1</b>	<b>27</b>	<b>1,168,978</b>

### **Age and Growth**

Biological data was collected from 204 CWT Chinook salmon to provide information on age, growth, and stocking-and-rearing history (Table 3). All of the CWT female Chinook salmon (N=172) returned as age 4 fish. No CWT female salmon returned as age 3 fish because no salmon were stocked in Lake Oahe in 2001. One hundred and seventy-two female salmon returned as age 4 fish mean length 802 mm (31.6 inches) and post-spawn mean weight 4,518g (9.95 lbs.) compared to mean length of 796 mm (31.3 inches) and post spawn weight 4,171g (9.19 lbs.) in 2002. In 2001 mean length of age 4 female salmon was 683 mm (26.9 inches) and mean weights were not available. Thirty-two CWT age 4 male Chinook returned to the station in 2003 with a mean length of 830 mm (32.7 inches) and weight 5,284 g (11.63lbs.). The ratio of male to female CWT salmon that returned to the station was 1:5.4 in 2003.

The number of male salmon that returned to WBS in 2003 was low compared to previous CWT tag male: to female ratios of 4.1:1 and 3.4:1 in 2000 and 2001. The large shift in the sex ratio was probably due a large number of male chinook salmon maturing, spawning and dying as 2 and 3 year old fish and therefore being absent from 2003 salmon run. The age composition of 2 and 3 year male CWT salmon have composed up to 100% of total CWT males returning to the station during previous spawns (Marrone and Stout 1995). No CWT male salmon returned as "jacks" or age 2 fish because no salmon were stocked in 2002. The 2003 run of Lake Oahe salmon produced the largest average length and weight of age 4 males and largest average length of age 4 female salmon since the station has been in operation. The largest average post-spawn mean weight of age 4 female salmon returned to the station in 1993 with a mean weight of (10.45 lbs.; Marrone and Stout 1993). The large average size of salmon in 2003 was likely attributed to a diet composed of rainbow smelt (*Osmerus mordax*), lake herring (*Coregonus artedii*), and gizzard shad (*Dorosoma cepedianum*).

**Table 3.** Age composition, length and weight of 204 coded-wire tagged Chinook salmon collected at Whitlocks Bay Spawning Station, fall 2003.

Age	Sex	Brood year	Number	Mean length (mm)	Range	Mean weight* (g)	Range
1	Male	2002	0	-----	-----	-----	-----
2	Male	2001	0	-----	-----	-----	-----
3	Male	2000	0	-----	-----	-----	-----
	Female		0				
4	Male	1999	32	830	630-907	5,284	2,253-7,491
	Female		172	805	654-930	4,518*	2,252-7,886

\*Female weight is post-spawn.

One of the coded wire tagged salmon was a fish stocked in Lake Sakakawea, North Dakota.

A breakdown of the total number, size at stocking and number of CWT salmon stocked in Lake Oahe from 1997 to the present can be found in Appendix F. In addition, South and North Dakota CWT codes and numbers of all tagged fish, from the start of tagging to the present, can be found in Appendices G and H. Detailed breakdowns of treatment groups, number of fish marked, and research purposes behind marking can be found in Appendix I. A complete summarization and analyses of CWT returns from 1987 - 1996 can be found in a report by Lott et al. (1997).

### Contribution of North Dakota Salmon

Of the 204 CWT Chinook salmon that returned to the WBS in 2003, one was from North Dakota. A percentage was determined using the known number of North Dakota fish tagged from each stock and calculating a percent return from the number (that returned to the spawning station) of North Dakota fish spawned at WBS. In 2003 it was estimated that 18 (i.e., approximately 4%) of the 459 salmon that

returned to the station were North Dakota fish. In 2002 it was estimated 22 (i.e., approximately 3%) of the 727 salmon that returned to the station were from North Dakota and in 2001 no coded wire tagged salmon returned to WBS. This estimate provides a very rough picture of the contribution of North Dakota salmon to the Chinook salmon spawning effort at Whitlock's Bay. No electrofishing was conducted in 2003 for Chinook salmon. North Dakota fish were not imprinted on Whitlock Bay therefore it was expected that they would have a higher incidence of straying into bays in an attempt to spawn. Straying North-and-South Dakota Chinook salmon were effectively electrofished in backbays in 1999 and 1998. By not electrofishing for salmon from 2000 thru 2003 it is assumed a large percentage of these straying fish would not be collected to spawn. This could partially explain the reason for the decreased or nonexistent contribution of North Dakota Chinook salmon to Lake Oahe spawn from 2000 thru 2003. However, water releases probably determined the presence or absence of North Dakota salmon in Lake Oahe. The large percentage of non-imprinted CWT North Dakota salmon in 1998 and 1999 are probably related to high water releases from Garrison Dam in 1996 and 1997 and these salmon likely contributed to the large number of fish that were straying into bays in an attempt spawn. This cannot be quantified because CWT tagged salmon captured electrofishing were not individually analyzed to determine the straying ratio of North Dakota versus South Dakota fish.

### **Tri-State Chinook Salmon Egg Take and Inter-State Assistance**

With few outside sources for disease free Chinook salmon eggs and large scale annual fluctuations in the number and quality of salmon eggs collected in Lake Oahe, Lake Sakakawea, and Fort Peck reservoir, tri-state cooperation is a necessity. This cooperation has come in the form of many things (i.e., sharing of CWT machines, research results, manpower, etc.) with the most obvious form of cooperation being the transfer of Chinook salmon eggs or fry/fingerlings among states when needed. A complete summary of South Dakota, North Dakota, and Montana Chinook salmon total egg take and inter-state egg, fry and fingerling transfers can be found in Appendix N. It is hoped that tri-state cooperation will continue and expand as the popularity of salmon fisheries on the prairie mainstem Missouri River reservoirs expands.




Figure 3. Chinook salmon returns to Whitlocks Bay Spawning Station compared to station intake water temperatures, Lake Oahe, SD, fall 2003.

Past stocking records of Chinook salmon for Lake Oahe are presented in Appendix B, C and E and a summary of past salmon returns and egg collection efforts can be found in Appendix E. An explanation of how the cost breakdown was completed for each category can be found in Appendix K.

**Table 4.** Costs of taking Chinook salmon eggs at Whitlocks Bay Spawning Station, 2003.

---

*Manpower/per diem/travel	\$7,122.00
*Utilities	2,495.00
**Miscellaneous-pump removal, hardware, etc	<u>11,103.00</u>
<b>Total Costs</b>	<b>\$20,720.00</b>
Number of eggs taken	1,168,978
Cost/1,000 eggs collected	\$ 17.73

---

\*Note previous reports used a value of \$20,000 for manpower/per diem/travel costs. An updated value is calculated for this category that more accurately reflects actual manpower/per diem/travel costs. Since 2000 the cost for the salmon eggs was computed from all expenses incurred during the entire year instead of just half year as in previous reports.

\*\*Costs associated with pump upgrade (construction of two individual intakes and submersible sleds, additional pipe and wire) were not included in the cost for the salmon eggs due to expenses that were part of the upgrade costs. The construction and delivery of 20 additional ladder sections were calculated into the miscellaneous cost. The ladder sections were necessary to operate the station due to low lake elevations.

### **Visitation and Tours**

The Whitlocks Bay Spawning Station was in operation from October 3 - October 21, 2003. Approximately 200 people visited the facility during that time. Personnel were available daily to answer questions, give tours and explain fisheries programs. The fish ladder was open for viewing everyday. Many radio, television, and newspaper interviews were provided to the news media. Special tours were given to two organized groups including schools and civic organizations.



### **Literature Cited**

- Hanten, R. and D. Stout 2000. 1999 Whitlocks Bay Spawning and Imprinting Station annual report. South Dakota Department of Game, Fish and Parks, Wildlife Division, Annual Report 00-02.
- Haskell, D. G. 1959. Trout growth in hatcheries. New York Fish and Game Journal 5:204-237.
- Johnson, B., R. Hanten, J. Lott, W. Nelson-Stastny 2000. Annual fish population and sport fish harvest surveys on Lake Oahe, South Dakota 1999. South Dakota Department of Game, Fish and Parks, Wildlife Division, Annual Report 01-14.
- Johnson, B., J. Lott, W. Nelson-Stastny and J. Riis 1997. Annual fish population and sport fish harvest surveys on Lake Oahe, South Dakota 1996. South Dakota Department of Game, Fish and Parks, Wildlife Division, Annual Report 97-15.
- Lott, J., R. Hanten, B. Johnson, W. Nelson-Stastny 1999. Annual fish population and sport fish harvest surveys on Lake Oahe, South Dakota 1999. South Dakota Department of Game, Fish and Parks, Wildlife Division, Annual Report 00-11.
- Lott, J., G. Marrone, D. Stout 1997. Influences of size-and-date at stocking, imprinting attempts and growth on initial survival, homing ability, maturation patterns and angler harvest of Chinook salmon in Lake Oahe, SD. South Dakota Department of Game, Fish and Parks, Wildlife Division, Report 97-20, Pierre.
- Marrone, G., M. and D. A. Stout 1995. 1995 Whitlocks Bay Spawning Station Annual Report. South Dakota Department of Game, Fish and Parks, Wildlife Division, Report, Pierre.
- Marrone, G., M. and D. A. Stout 1993. 1993 Whitlocks Bay Spawning Station Annual Report. South Dakota Department of Game, Fish and Parks, Wildlife Division, Report, Pierre.
- Scott, W., B. and E. J. Crossman. 1973. Freshwater fishes of Canada. Fisheries Research Board of Canada, Ottawa.

Appendix A. Rainbow trout stocked in Lake Oahe, 1996 - 2003.

Date	Number	Strain	No./lb.	Mark	Treatment	Location
04/15/96	45,298	O	11.5	None	None	WB
04/16/96	18,768	O	8.1	CW	None	WB
04/23/96	20,582	O	10.1	None	None	WB
04/25/96	10,150	O	9.9	None	None	WB
11/04/96	30,327	O	88.2	None	None	WB
11/04/96	20,534	O	56.5	CW	None	WB
04/16/97	80,480	O	11.3	None	None	WB
04/18/97	18,029	O	8.5	CW	None	WB
11/06/97	14,953	O	64.8	CW	None	WB
11/06/97	5,496	O	134.1	None	None	WB
04/13/98	29,162	O	13.65	CW	None	WB
04/20/98	62,157	O	15.38	None	None	WB
11/24/98	18,075	O	17.1	None	None	WB
11/24/98	29,206	O	17.1	CW	None	WB
04/19/99	28,742	O	9.5	None	None	WB
04/20/99	28,131	O	9.11	CW	None	WB
09/09/99	19,885	M	19.56	None	None	WB
09/14/99	17,776	M	16.97	None	None	WB
9/23/99	43,353	M	18.37	None	None	WB
10/27/99	18,669	O	14.43	CW	None	WB
11/03/99	3,331	O	20.46	None	None	WB
11/03/99	9,806	O	15.91	None	None	WB
4/12/00	15,172	O	6.05	CW	Hand Fed	EWB
4/12/00	15,408	O	5.93	CW	Auto Feeder	EWB
4/17/00	2,879	O	7.17	None	None	EWB
7/19/00	67,562	O	47.28	None	None	EWB
2001	0	-	-	-		-
2002	0	-	-	-	-	
2003	0	-	-	-		-

CW = coded-wire tagged and adipose fin clipped; AD = adipose fin clipped fish

Appendix B. Numbers of coldwater species stocked in Lake Oahe, 1983 - 2003

<b>Year</b>	<b>Brown trout</b>	<b>Chinook salmon</b>	<b>Lake trout</b>	<b>Rainbow trout</b>	<b>Lake herring</b>
1983		790,150	7,779	131,395	
1984	18,000	784,550	8,560	279,395	470,000
1985	69,000	845,542	4,750	85,635	-----
1986	52,900	811,665	-----	146,229	
1987	93,700	1,005,054	-----	50,000	
1988	77,167	1,061,535		41,270	200
1989	50,000	217,037	-----	19,150	
1990	50,500	66,385		32,370	9,388,500
1991	-----	249,478	-----	18,410	10,995,500
1992	-----	219,000	-----	27,190	11,416,000
1993	-----	275,055		74,707	-----
1994		298,759		81,365	-----
1995		363,787		96,002	
1996	-----	418,478	-----	145,659	
1997	-----	396,660	-----	118,958	
1998	-----	234,101	-----	131,245	
1999	-----	104,976	-----	213,506	
2000		49,494		101,021	-----
2001	-----		-----		
2002	-----	-----	-----		
2003		17,259	-----	-----	-----

Appendix C. Pounds and kilograms of coldwater species stocked in Lake Oahe, 1982 – 2003.

Year	Brown trout		Chinook salmon		Lake trout		Rainbow trout		Steelhead trout		Total	
	(lbs.)	(kg.)	(lbs.)	(kg.)	(lbs.)	(kg.)	(lbs.)	(kg.)	(lbs.)	(kg)	(lbs.)	(kg.)
1982	----	----	2,169	984	1,373	623	1,988	902	40	18	5,570	2,527
1983	----	----	8,889	4,032	1,903	863	1,532	695	----	----	12,324	5,590
1984	1,385	632	11,266	5,110	2,094	950	12,541	5,689	----	----	27,286	12,377
1985	4,182	1,908	11,882	5,390	1,041	472	1,691	767	430	195	19,226	8,721
1986	5,290	2,414	11,132	5,049	----	----	7,901	3,584	6,256	2,838	30,579	13,871
1987	4,462	2,036	15,754	7,146	----	----	----	----	12,500	5,670	32,716	14,840
1988	5,599	2,555	13,810	6,264	----	----	----	----	8,351	3,788	27,760	12,592
1989	3,378	1,541	9,299	4,218	----	----	----	----	2,217	1,006	14,894	6,756
1990	3,389	1,546	2,417	1,096	----	----	8,093	3,671	----	----	13,899	6,305
1991	----	----	8,473	3,843	----	----	3,610	1,637	----	----	12,083	5,481
1992	----	----	5,300	2,404	----	----	3,153	1,430	----	----	8,453	3,834
1993	----	----	7,348	3,333	----	----	5,257	2,385	----	----	12,605	5,718
1994	----	----	8,169	3,705	----	----	6,732	3,054	----	----	14,901	6,759
1995	----	----	9,922	4,501	----	----	9,973	4,524	----	----	19,895	9,024
1996	----	----	10,893	4,941	----	----	10,022	4,546	----	----	20,915	9,487
1997	----	----	5,053	2,292	----	----	9,492	4,306	----	----	14,545	6,598
1998	----	----	9,428	4,277	----	----	8,942	4,056	----	----	18,370	8,333
1999	----	----	7,608	3,451	----	----	14,647	6,644	----	----	22,255	10,095
2000	----	----	6,240	2,830	----	----	6,937	3,147	----	----	13,177	5,977
2001	----	----	----	----	----	----	----	----	----	----	----	----
2002	----	----	----	----	----	----	----	----	----	----	----	----
2003	----	----	1,121	508	----	----	----	----	----	----	----	----
<b>Total</b>	27,685	12,633	165,052	74,868	6,411	2,908	112,511	51,035	29,794	13,515	341,453	154,883
<b>Average</b>	3,955	1,805	8,309	3,769	1,603	727	7,032	3,190	4,966	2,252	17,971	8,152

Totals and averages were calculated from years when fish were stocked.

Appendix D. Summary of rainbow trout return and egg collection at Whitlocks Bay Spawning Station, 1984 -2003.

Year	Total rainbows returned	Number of females spawned	Number of taken	Average number of eggs per female	percent egg eye-up
1984				----	
1985	----				
1986	----	----			
1987	50	none	----	----	----
1988	57	22	36,915	1,678	----
1989	8	none		----	
1990	7	none		----	
1991	126	39	52,675	1,351	----
1992	296	127	245,660	1,934	
1993	276	121	319,500	2,640	75
1994	641	170	376,620	2,215	81
1995	340	131	469,085	3,581	71
1996	323	147	411,690	2,801	78
1997	203	95	260,302	2,740	79
1998	111	46	131,245	2,853	60
1999	74	30	70,870	2,362	80
2000	138	47	120,300	2,560	69
2001	----		----	----	----
2002	----		----	----	----
2003					
Total	2650	975	2,494,862		----
Average	189	89	226,806	2,429	74

Appendix E. Summary of salmon return and *egg* collection at Whitlocks Bay Station\_ 1984-2003.

<b>Year</b>	<b>Total salmon returned</b>	<b>Number of females spawned</b>	<b>Number of good eggs</b>	<b>Average number of eggs/female</b>	<b>Percent egg eye-up</b>
1984	175	29	81,340	2,800	-----
1985	435	119	364,900	3,066	-----
1986	484	61	186,746	3,061	
1987	1,034	187	871,137	4,659	-----
1988	1,320	270	667,796	2,473	
1989	1,256 <sup>a</sup>	227	418,160	1,842	37
1990	1,107	409	787,708	1,926	40
1991	1,343	193	453,864	2,352	60
1992	690 <sup>b</sup>	319	992,630	3,112	40
1993	1,354	449	1,744,100	3,884	53
1994	2,749	408	1,542,180	3,780	44
1995	3,664	527	1,969,162	3,737	48
1996	1,133	355	1,054,841	2,971	74
1997	673 <sup>c</sup>	193	558,945	2,896	53
1998	422 <sup>d</sup>	211	498,100	2,361	48
1999	328 <sup>e</sup>	124	316,862	2,555	70
2000	283	74	155,700	2,100	49
2001	468	67	153,264	2,286	31
2002	727	189 <sup>f</sup>	724,380 <sup>f</sup>	3,833 <sup>f</sup>	57 <sup>f</sup>
2003	459	284	1,168,978	4,116	48
Total	20,104	4,695	14,710,793	59,810	-
Average	1,005	235	735,540	2,991	50

<sup>a</sup> includes 71 females and 121 males from Spring Creek, Sutton Bay and Whitlocks Bay.

<sup>b</sup> includes 307 females from East Shore, Cow/Spring Creek and Cheyenne Creek

<sup>c</sup> includes 124 female and 33 male from Co/Spring/Okobojo Crks., Son Ba, Cheyenne Creek

<sup>d</sup> includes 131 females from Cow/Spring Creeks, West Shore, Sutton and Whitlocks Bay

<sup>e</sup> includes 76 females from Cow/Springs Creeks and Sutton and Whitlocks Bay

<sup>f</sup> does not include 134 females eggs stripped and fertilized at hatchery (467,992 eggs collected 11% eye-up

Appendix F. Chinook salmon stocked by number, size and mark in Lake Oahe, 1998 - 2003.

Date	Number	Strain	Hatchery	No./lb.	Mark	Fin clips	Location
04/27/98	35,070	O	BD	105	None	None	WB
05/27/98	26,075	O	BD	35.0	None	None	WB
06/02/98	28,822	O	MN	34.5	CW	AD	WB
06/02/98	97,292	O	MN	37.7	None	None	WB
09/23/98	18,461	O	MN	9.4	CW	AD	WB
09/30/98	28,381	O	MN	9.6	None	None	WB
06/02/99	15,745	O	MN	40.24	CW	AD	WB
06/02/99	14,519	O	MN	43.13	CW	AD	WB
06/10/99	24,508	O	MN	31.9	None	None	WB
09/29/99	19,371	O	MN	8.2	CW	AD	WB
09/29/99	30,873	O	MN	8.2	None	None	WB
07/19/00	7,458	O	MN	16.46	None	None	WB
09/20/00	15,000	O	MN	8.46	CW	AD	WB
09/26/00	19,494	O	MN	8.58	None	None	WB
09/26/00	15,000	O	MN	8.61	CW	AD	WB
2(X)1		----	----				----
2002	----	----	----	----	----		
06/02/03	5,688	----	MN	34.76	CW	AD	WB
06/02/03	4,403	O	MN	33.64	CW	AD	WB
10/27/03	4,368	O	MN	8.43	CW	AD	WB
11/18/03	2,800	O	CS	9.10	CW	None	WB

CW = coded-wire tagged and adipose fin clipped; AD = adipose fin clipped fish Hatchery codes BD, MN and CS is Blue Dog, McNenny and Cleghorn Springs State Fish Hatcheries

Appendix G. South Dakota Chinook salmon coded-wire-tag data 1989 -2003.

Year stocked	Brood year	Description	Number tagged	Number stocked	Tag retention (percent)	Datal	Data2
1990	1989	WBS/Fall Stock (8/lb.)	10,300			17	19
1990	1989	WBS/(53/lb.)	29,019	-----		19	26,27
1991	1989	WBS (2.9/lb.)	11,990	-----		19	24
1991	1989	OATW (2.9/lb.)	5,254	-----		19	25
1991	1990	WWB Direct Stock (45/lb.)	19,242	-----		20	28,29,30
1991	1990	WBS Stock	17,850	-----		20	31,32,33,34
1992	1991	WWB (30/lb.)	30,193	-----	89.5%	21	1,2
1992	1991	WWB (runts)	2,623	-----	86.0%	21	3
1993	1992	WWB (34/lb.)	22,959	-----	92.5%	22	1,2,3
1993	1992	WWB (51/lb.)	18,430	-----	80.0%	22	4,5,6,7
1994	1993	WWB (29.6/lb.)	47,973	210,656	82.8%	23	1,2,3,4,5,6
1995	1994	WWB (27.9/lb.)	48,469	212,999	98.0%	24	1,2,3,4,5,6
1996	1995	WWB (32.5/lb.)	46,394	237,217	86.0%	25	1,2,3,4,5,6
1997	1996	WWB (38.9/lb.)	50,428	243,278	80.5%	26	1,2,3,4,5,6
1998	1997	WWB (34.5/lb.)	28,822	126,144	82.1%	27	1
1998	1997	WWB (9.41/lb.)	19,315	47,696		27	2
1999	1998	WWB (43.13/lb.)	14,519	54,772	99%	28	1
1999	1998	WWB (40.24/lb.)	15,745	54,772	97.1%	28	2
1999	1998	WWB (8/lb.)	9,686	50,204	98.6%	28	3
1999	1998	WWB (8.12/lb.)	9,685	50,204	98.3%	28	4
2000	1999	WWB (8.61lb.)	14,850	24,597	99%	29	1
2000	1999	WWB (8.46/lb.)	14,835	24,582	98.9%	29	2
2001	-----	-----	-----	-----	-----	-----	-----
2002	-----	-----	-----	-----	-----	-----	-----
2003	2002	WWB (33.64/lb.)	4,403	4,403	87.5%	25,28	8,6
2003	2002	WWB (34.76lb.)	5,688	5,688	39.2%	22	6,8
2003	2002	WWB (8/lb.)	4,368	4,368	25.1%	22	9



Appendix H. North Dakota Chinook salmon coded-wire-tag data 1993 -2003.

Year stocked	Brood year	Description	Number tagged	Number stocked	Data1	Data2
1994	1993	Cage reared (28/lb.)	15,000	75,000	28	7
1994	1993	GDNFH (37/lb.)	15,000	204,500	28	1
1994	1993	Cage reared (34/lb.)	15,000	123,000	28	3
1995	1994	Cage reared (36/lb.)	13.5(X)	74,000	24	2
1995	1994	GDNFH (8/lb.)	13.5(X)	23,665	24	6
1995	1994	Cage reared (35/lb.)	13,500	130,000	24	14
1995	1994	GDNFH (33/lb.)	13,500	137,247	30	8
1996	1995	Cage reared (49/lb.)	13,000	120,000	30	4
1996	1995	Cage reared (44/lb.)	13,000	79,000	30	5
1996	1995	GDNFH (39/lb.)	13,000	170,000	30	10
1996	1995	GDNFH (44/lb.)	13,000	120,011	30	12
1997	1996	Cage reared (82/lb.)	52,900	38,059	31	1
1998	1997	Small Cage reared (51/lb.)	18,327	70,000	1	2
1998	1997	Large Cage reared (28/lb.)	18,327	69,500	1	5
1998	1997	GDNFH (41/lb.)	18,327	86,482	1	3
1999	1998	GDNFH (96/lb.)	15.9(X)	131,325	60	1
1999	1998	GDNFH (65/lb.)	15.9(X)	92,002	60	2
2000	1999	GDNFH (60/lb.)	15,000	250,000	60	3
2000	1999	GDNFH (90/lb.)	15,000	250,000	60	4
2001	2000	GDNFH (62/lb.)	11,386	255,276	1	72
2001	2000	GDNFH (62/lb.)	602		45	58
2001	2000	GDNFH (62/lb.)	2,012		24	14
2001	2000	GDNFH (106/lb.)	11,248	205,300	1	71
2001	2000	GDNFH (106/lb.)	3,752		30	08
2002	2001	Cage-Rodeo (25/lb.)	14,000	110,007	2	2
2002	2001	Cage-Rodeo (50/lb.)	14,000	98,283	2	1
2002	2001	GDNFH-Rodeo (50/lb.)	14,000	250,554	2	3
2003	2002	GDNFH-Rodeo (27/lb.)	15,000	97,500	2	12
2003	2002	GDNFH-Rodeo (45/lb.)	15,000	202,965	2	11

Appendix I. South Dakota rainbow trout coded-wire tag data 1991 -2003.

Year stocked	Brood year	Description	Number tagged	Number stocked	Tag retention (percent)	Datal	Data2
1992	1991	WWB (5.68/lb.)	4,135		981/6	20	3536,37
1992	1991	WWB (10.07/lb)	8,115		83%	20	35,36,37
1992	1991	WWB (6.38/lb.)	5,270			20	35,36,37
1992	1991	WWB (12.2/lb.)	9,670			20	3536,37
1994	1993	WWB (9.51/16.), CW	19,440	81,365	93.8%	21	4,5
1995	1994	WWB (9.82/lb.), CN	22,891	100,002	93.9%	23	7,8,9,10
1995	1994	WWB (53.3/lb.), CN	19,655	34,655	93.9%	23	7,8,9,10
1996	1995	WWB (56.5/16.), CN	18,768	50,861	94.0%	24	7,8
1996	1995	EWB (8.12/lb.), CN	19,655	94,798	85.6%	24	9,10
1997	1996	WWB (64.8/16.), CN	14,953	20,089	82.7%	25	7,8,9
1997	1996	WWB (8.54/lb.), CN	18,029	98,509	84.5%	23	6,9,10
1997	1996	WWB (8.54/lb.), CN			84.5%	17	17,20,21
1997	1996	WWB (8.54/lb.), CN			84.5%	20	33
1998	1997	WWB (13.65/lb.), CW	29,162	91,319	79.4%	26	6,7,8
1999	1998	WWB (9.11/lb), CW	28,131	56,873	89.9%	27	6
1999	1999	WWB (14.43/lb.), CW	18,669	41,654	%	28	8
2000	1999	WWB (5.93/lb.); CW	15,408	15,408	74.1%	28	7
2000	1999	WWB (6.05/lb.); CW	15,172	15,172	62.5%	28	6
2001	2000	None	----	----	----	----	----
2002	2001	None	----	----	----	----	-
2003	2002	None	----	----	----	----	----

CW = coded-wire tagged and adipose fin clipped; CN = coded-wire tagged no clip

Appendix J. Number of coded-wire-tagged rainbow trout stocked in Lake Oahe, 1987 - 2003. Number stocked is corrected for tag retention.

Date stocked	Location	Size (number/lb.)	Code	Number stocked	Treatment	Purpose
4/12/2000	East Whitlock Bay	6.05	28-7	15,172	Hand fed	Determine how feeding practices influence
4/12/2000	East Whitlock Bay	5.93	28-6	15,408	Auto Fed	post-stocking salmonid survival
2001	-----	----	----	Suspended		Study is suspended due to prey shortage when rainbow trout stocking resumes the study will continue
2002	-----	----	----	Suspended		Study is suspended due to prey shortage when rainbow trout stocking resumes the study will continue
2003	-----	----	----	Suspended		Study is suspended due to prey shortage when rainbow trout stocking resumes the study will continue

Appendix K. Number of coded-wire-tagged Chinook salmon stocked in Lake Oahe, 1987 - 2003.  
Number stocked is corrected for tag retention

Date stocked	Location	Size (number/lb.)	Number stocked	Treatment	Purpose
04/15/87	Whitlocks Bay	70.0	29,602	21 day morpholine imprint at Blue Dog and held 14 days at Whitlocks station.	Evaluate chemical imprinting and stocking size.
04/15/87	Whitlocks Bay	70.0	29,910	Held at Whitlocks station 14 days.	
05/18/87	Whitlocks Bay	32.0	19,876	Held at Whitlocks station 14 days.	
05/18/87	Whitlocks Bay	32.0	20,188	21day morpholine imprint at Blue Dog And held 14 day at Whitlocks Station	
05/12/88	Whitlocks Bay	64.0	64,508	Held at Whitlocks station 10 days.	Evaluate stocking size and
05/31/88	Whitlocks Bay	30.0	53,815	Held at Whitlocks station 14 days.	identify Oahe stock.
04/12/89	West Shore	3.3	6,235	None	Evaluating yearling stocking size
04/13/89	Whitlocks Bay	3.3	6,250	Held at Whitlocks station 4 days.	and stocking location. Evaluate
05/25/89	Whitlocks Bay	43.8	28,999	Held at Whitlocks station 10 days.	adult return rate of smolts held at
06/05/89	Whitlocks Bay	34.4	30,164	None- directly stocked into Whitlocks Bay from hatchery	Whitlocks station vs. those stocked directly.
06/08/90	Whitlocks Bay	53.0	29,019	Held at Whitlocks station 25 days	Evaluate stock size and identify
11/01/90	Whitlocks Bay	8.0	7,704	Held at Whitlocks station 15 days.	known age fish. Evaluate fall
04/24/91	Whitlocks Bay	2.9	9,232	None - direct stock	stock and identify know age fish.
03/08/91	Oahe Tailwaters	3.0	5,254	None	Evaluate stock size and identify known age and hatchery source. evaluate Oahe Tailwater stocking.
05/15/91	Whitlocks Bay	45.0	17,850	Held at Whitlocks station 14 days.	Compare adults return rate from
05/16/91	Whitlocks Bay	46.2	19,242	None - direct stock.	smolts held and stocked from Whitlocks station.

Appendix K. Number of coded-wire-tagged Chinook salmon stocked in Lake Oahe, 1987 - 2003.

Number stocked is corrected for tag retention (continued).

Date stocked	Location	Size (No./lb.)	Code	Number stocked	Treatment	Purpose
05/27/92	Whitlocks Bay	29.5		30,193	None - direct stock.	Identify known age fish.
05/27/92	Whitlocks Bay	60.3		2,623	None - direct stock	determine stocking success and survival of "runs".
05/25/93	Whitlocks Bay	33.9		22,959	None - direct stock	Identify known age fish. Compare
05/25/93	Whitlocks Bay	50.6		18,430	None - direct stock	stocking sizes of 34/lb. with 51/lb.
05/23/94	Whitlocks Bay	29.6		47,973	None - direct stock	Identify known age fish.
05/30/95	Whitlocks Bay	27.9		48,469	None - direct stock	Identify known age fish.
05/28/96	Whitlocks Bay	32.5		46,397	None - direct stock	Identify known age fish.
05/27/97	Whitlocks Bay	38.9		50,428	None - direct stock	Identify known age fish.
06/02/98	Whitlocks Bay	34.5		28,822	None - direct stock	Identify known age fish.
09/23/98	Whitlocks Bay	9.4		18,461	None - direct stock	
06/02/99	Whitlocks Bay	40.2	28-2	15,745	Inside circular tank rearing density <10 kg/m <sup>3</sup>	Determine how rearing density
06/02/99	Whitlocks Bay	43.1	28-1	14,519	Inside circular tank rearing density >20 kg/m <sup>3</sup>	and water flows influence
09/29/99	Whitlocks Bay	8.0	28-3	9,686	Reared in covered raceway at low flow (200gpm)	post-stocking salmonid survival
09/29/99	Whitlocks Bay	8.1	28-4	9,685	Reared in covered raceway at high flow (>300gpm)	
09/26/00	Whitlocks Bay	8.46	29-2	14,835	Reared in covered raceway at low flow (>100gpm)	Determine how water flows influence
09/26/00	Whitlocks Bay	8.61	29-1	14,850	Reared in covered raceway at high flow (>250gpm)	post-stocking salmonid survival
2001	-----				Suspended	Study was suspend until FCS stockings
2002	-----				Suspended	continue due to a prey shortage
06/02/03	Whitlocks Bay	33.64	22-6,8	3,853	Inside circular tank rearing density <10 kg/m <sup>3</sup>	Replicate 1999
06/02/03	Whitlocks Bay	34.76	25-8,28-6	2,230	Inside circular tank rearing density >20 kg/m <sup>3</sup>	Replicate 1999
10/27/03	Whitlocks Bay	8.43	22-9	1,096	Growth/survival data	Replicate

\* Code designation not determined yet.

Appendix L. Percent recovery of coded-wire tags returned by anglers and recovered at Whitlocks Bay Spawning Station in Lake Oahe, 1988-2003.

Year stocked	Brood year	Description	Total number stocked	% Tag retention	Corrected #tagged fish	Total No. tagged fish return (fisherman)	Total No. tagged fish return (station)	% of tagged FCS returned to				
								Fisherman	Station	Total	Data 1	Data 2
1988	1987	Large (30/lb.)	-----	100%	53,815	183	125	0.34	0.23	0.57	17	1,2,3,4,5,6,7,8
1988	1987	Small (64/lb.)	-----	100%	64,508	90	49	0.14	0.08	0.22	17	9,10,11,12,14,15
1989	1987	Oahe Dam (3.3/lb.)	-----	100%	6,235	44	2	0.71	0.03	0.74	17	16
1989	1987	Whitlocks (3.3 lb.)	-----	100%	6,250	28	67	0.45	1.07	1.52	17	18
1989	1988	Whitlock Stock (44/lb.)	-----	92.60%	31,312	66	93	0.21	0.30	0.51	18	22
1989	1988	Direct Stock (34.5/lb.)	-----	100%	30,164	83	146	0.28	0.48	0.76	18	23
1990	1989	WBS/Fall Stock (8/lb.)	-----	74.80%	10,300	93	108	0.90	1.05	1.95	17	19
1990	1989	WBS/(53/lb.)	-----	100%	29,019	114	106	0.39	0.37	0.76	19	26,27
1991	1989	WBS (2.9/lb.)	-----	76.99%	11,990	177	381	1.48	3.18	4.65	19	24
1991	1990	WWB Direct Stock (45/lb.)	-----	100%	19,242	107	106	0.56	0.55	1.11	20	28,29,30
1991	1990	WBS Stock	-----	100%	17,850	82	114	0.46	0.64	1.10	20	31,32,33,34
1992	1991	WWB (30/lb.)	-----	89.50%	30,193	186	199	0.62	0.66	1.28	21	1,2
1992	1991	WWB (runs; 60.5/lb.)	-----	86.00%	2,623	2	15	0.08	0.57	0.65	21	3
1993	1992	WWB (34/lb.; Direct stock)	-----	92.50%	22,959	169	220	0.74	0.96	1.69	22	1,2,3
1993	1992	WWB (51/lb.)	-----	80.00%	18,430	104	108	0.56	0.59	1.15	22	4,5,6,7
1994	1993	WWB (29.6/lb.)	210,656	82.80%	47,973	269	379	0.56	0.79	1.35	23	1,2,3,4,5,6
1995	1994	WWB (27.9/lb.)	212,999	98.00%	48,469	65	87	0.13	0.18	0.31	24	1,2,3,4,5,6
1996	1995	WWB (32.5/lb.)	237,217	86.00%	46,394	10	18	0.02	0.04	0.06	25	1,2,3,4,5,6
1997	1996	WWB (38.9/lb.)	243,278	80.50%	50,428	23	96	0.05	0.19	0.24	26	1,2,3,4,5,6
1998	1997	WWB (34.5/lb.)	126,144	82.10%	28,822	7	14	0.02	0.05	0.07	27	1
1998	1997	WWB (9.41/lb.)	47,696		19,315	4	28	0.02	0.14	0.17	27	2
1999	1998	WWB (43.13/lb.)	54,772	99.00%	14,519	-	3	0.00	0.02	0.02	28	1
1999	1998	WWB (40.24/lb.)	54,772	97.10%	15,745	2	26	0.01	0.17	0.18	28	2
1999	1998	WWB (8/lb.)	50,204	98.60%	9,686	2	59	0.02	0.61	0.63	28	3
1999	1998	WWB (8.12/lb.)	50,204	98.30%	9,685	2	55	0.02	0.57	0.59	28	4
2000	1999	WWB (8.61/lb.)	24,597	99.00%	14,850	4	180	0.03	1.21	1.24	29	1
2000	1999	WWB (8.46/lb.)	24,582	98.90%	14,835	-	116	0.00	0.78	0.78	29	2
2001	2000	None										
2002	2001	None										
2003	2002	WWB (33.7/lb.)	5,688	39.20%	2,230						25, 28	8, 6
2003	2002	WWB (33.6/lb.)	4,403	87.50%	3,853						22	6, 8
2003	2002	WWB (8.4/lb.)	4,368	25.1%	1,096						22	9

Portions in gray are still being created

## Appendix M. Salmon and trout cost breakdown explanation for each category in spring and fall budgets

In the 2003 Whitlocks Report the egg take cost was calculated for the Chinook salmon spawn only. All expenses incurred in 2003 were included in the salmon egg costs. Prior to 2003, cost breakdowns for the spring rainbow trout spawn and fall Chinook salmon spawn were divided in half. Spring trout spawning costs cover all spawning station operation costs from January 1<sup>st</sup> to July 15<sup>th</sup> and salmon spawning costs cover all spawning station operation costs from July 16<sup>th</sup> to the end of the year.

Costs associated with manpower, per diem and travel to and from the spawning station are factored into the overall spawning costs. A considerably larger portion of this budget is weighted to the fall salmon spawn that requires more people to complete.

Utilities include electricity demand and use charges, and telephone costs. The miscellaneous category includes office and hardware supplies; building and grounds maintenance costs; equipment rental (e.g., CO<sub>2</sub> and Oz bottle rental); recreational supplies; vehicle maintenance and repair; propellants/oils/lubricants; rock, sand, gravel etc. Costs associated with pump upgrade (construction of two individual intakes and submersible sleds, additional pipe and wire) were not included in the cost for the salmon eggs due to expenses that were part of the upgrade costs. The construction and delivery of 20 additional ladder sections were calculated into the miscellaneous cost. The ladder sections were necessary to operate the station due to low lake elevations.

Appendix N. Calculation of estimated contribution of North Dakota Chinook salmon to the spawn of Lake Oahe, South Dakota 2003.

ID code	State	Year stocked	Brood year	Stocking size /lb.	ND FCS No. CWT	No. fish captured	% CWT returned	Total No. stocked	Est. No. of ND FCS
	ND	2001	2000	106	11,248	1	0.00889	205,300	18
<b>TOTALS</b>					11,248	1			18
Estimated % of ND									
Total FCS Est. No. of FCS collected during									
collected ND FCS during SD Spawn									
459	18	3.98%							

Note: Data used in calculations from Jason Lee, North Dakota Fish and Game Department. CWTTREAT.DBF - Table Records



Appendix O. Summary of South Dakota, North Dakota and Montana Chinook salmon total egg take, distribution, or inter-state assistance.

Year	No. of eggs eggs taken in SD	No. of eggs eggs taken in ND	No. of eggs eggs taken in MT	No. of excess eggs given to ND from SD	No. of excess eggs given to MT from SD	No. of excess eggs given to SD from ND	No. of excess eggs given to MT from ND
1980		90,000					
1981	100,000	400,000					
1982		Unknown					
1983		N/A					250,000
1984	81,340	395,000					
1985	364,900	379,700					
1986	186,746	479,600					177,413
1987	871,137	2,403,000					
1988	667,796	933,234					55,940
1989	418,160	739,350					
1990	787,708	1,574,269					101,210
1991	453,864	654,280					
1992	992,630	1,087,500			112,000		
1993	1,744,100	392,000		677,200			
1994	1,542,180	381,700	0 <sup>a</sup>	502,500	50,000		37,000
1995	1,969,162	547,650		849,000	240,000		
1996	1,054,841	1,570,130	390,000	228,000			
1997	558,945	1,126,406				50,000	
1998	498,100	749,116	6,000		49,561		
1999	316,862	3,038,219	141,000		193,000 <sup>b</sup>		250,000
2000	155,700	2,203,638			54,925		344,393
2001	153,264	2,400,322	0 <sup>a</sup>		43,150		600,000
2002	1,192,372	2,005,380	64,000		350,800 <sup>c</sup>		
2003	1,162,372	644,242	231,645	311,200	186,366		
<b>Total</b>	15,272,179	24,194,736	832,645	2,567,900	1,279,802	50,000	1,357,620
<b>Average</b>	727,247	1,099,761	118,949	513,580	142,200	50,000	169,703

<sup>a</sup> Zero indicates attempt was made to collect eggs

<sup>b</sup> Includes 100,000 eggs from collected from Cleghorn Springs Hatchery brood stock

<sup>c</sup> Includes 2000,000 fingerling raised for MT

Appendix P. Length at selected number per pound values for rainbow trout and Chinook salmon.

Length at selected number per pound values for rainbow trout

**Note:** These values come from condition factor averaged from  
Thousand of hatchery fish measured and weighed and are to be considered with  
A grain of salt for wild fish.

Condition factor (C)	Number per lb.	Individual weight lb.	Length (inches)	C is based on English measurements (pounds and inches)	
0.0004	2500.00	0.0004	1.0		
0.0(X)4	740.74	0.0014	1.5	Excel equation to calculate length from a known number/pound.	
0.0004	312.50	0.0032	2.0	=IOA((LOG(I000/#LB.)-LOG(Cond Factor* 1000))/3)	
0.0004	160.00	0.0063	2.5	<b>To calculate length in inches enter number/lb here</b>	
0.0004	92.59	0.0108	3.0	<b>and condition factor (C) here</b> 0.0004	
0.0004	58.31	0.0172	3.5	<b>length in inches =</b> 6.30	
0.0004	39.06	0.0256	4.0		
0.0004	27.43	0.0365	4.5		
0.0004	20.00	0.0500	5.0		
0.0004	15.03	0.0666	5.5		
0.0004	11.57	0.0864	6.0	Excel equation to calculate individual fish weight from a known Length	
0.0004	9.10	0.1099	6.5	<b>=Cond. Factor*(LengthA3)</b>	
0.0004	7.29	0.1372	7.0	<b>To calculate number/lb enter length (inches) here</b>	
0.0004	5.93	0.1688	7.5	<b>and condition factor (C) here</b> 0.0004	
0.0004	4.88	0.2048	8.0	<b>Weight per fish =</b> 0.1000	
0.0004	4.07	0.2457	8.5	<b>Number per pound =</b> 9.9981	
0.0004	3.43	0.2916	9.0		
0.0004	2.92	0.3430	9.5		
0.0004	2.50	0.4000	10.0		

Length at selected number per pound values for Chinook salmon

**note:** These values come from condition factor averaged from  
thousand of hatchery fish measured and weighed and are to be considered with  
a grain of salt for wild fish.

Condition factor	Number per lb.	Individual weight lb.	Length (inches)	Guide to selecting condition factor (C) for various fish species	
0.0003	3333.33	0.0003	1.0		
0.0003	987.65	0.0010	1.5		
0.0003	416.67	0.0024	2.0		
0.0003	213.33	0.0047	2.5	<b>Condition factor (C)</b>	<b>Species</b>
0.0003	123.46	0.0081	3.0		
0.0003	77.75	0.0129	3.5		
0.0003	52.08	0.0192	4.0		
0.0003	36.58	0.0273	4.5		
0.0003	26.67	0.0375	5.0		
0.0003	20.04	0.0499	5.5		
0.0003	15.43	0.0648	6.0		
0.0003	12.14	0.0824	6.5		
0.0003	9.72	0.1029	7.0		
0.0003	7.90	0.1266	7.5		
0.0003	6.51	0.1536	8.0		
0.0003	5.43	0.1842	8.5		
0.0003	4.57	0.2187	9.0		
0.0003	3.89	0.2572	9.5		
0.0003	3.33	0.3000	10.0		

Haskell, David G. 1959. Trout growth in hatcheries. New York Fish and Game Journal, Vol. 5, no. 2, pp. 204 - 237.

**Appendix Q.** Chinook salmon carcass contract results for 2003.

Date	Number of salmon received	Pounds of fish received	Bid price/lb	Total
Oct. 8, 2003	87	972	<b>\$0.20</b>	<b>\$ 194.40</b>
Oct. 14, 2003		916	<b>\$0.20</b>	<b>\$ 183.20</b>
Oct. 21 2002	141	1 384	<b>\$0.20</b>	<b>\$276.80</b>
<b>Total:</b>	228	3,272		<b>\$654.40</b>
<b>2003 cost for salmon carcasses:</b>				<b>\$654.40</b>

**Number tally does not include number of fish from October 14th  
date**

Note additional weight over 7,000 pounds rate changes to  
10 cents/pound.